

1                   **Accounting for demand and supply of the Biosphere's regenerative capacity:**  
2                   **the National Footprint Accounts' underlying methodology and framework**  
3  
4  
5

6                   Michael Borucke<sup>1</sup>, David Moore<sup>2</sup>, Gemma Cranston<sup>3</sup>, Kyle Gracey<sup>1</sup>, Katsunori Iha<sup>1</sup>, Joy Larson<sup>1</sup>, Elias  
7                   Lazarus<sup>1</sup>, Juan Carlos Morales<sup>1</sup>, Mathis Wackernagel<sup>1</sup>, Alessandro Galli<sup>2,\*</sup>  
8  
9

10  
11  
12                  <sup>1</sup> Global Footprint Network, 312 Clay Street, Oakland, CA, 94607-3510 USA  
13

14                  <sup>2</sup> Global Footprint Network, International Environment House 2, 7-9 Chemin de Balexert, 1219 Geneva  
15                  - Switzerland  
16

17                  <sup>3</sup>University of Cambridge, Programme for Sustainability Leadership, UK  
18

19                  \*Corresponding author:  
20

21                  Alessandro Galli, Ph.D.  
22                  Global Footprint Network,  
23                  International Environment House 2,  
24                  7-9 Chemin de Balexert,  
25                  1219 Geneva - Switzerland  
26                  Tel: +41 22 797 41 10  
27                  Mobile: +39 346 6760884  
28                  e-mail: [alessandro@footprintnetwork.org](mailto:alessandro@footprintnetwork.org)  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49                  NOTICE: This is the author's version of a work that was accepted for publication in *Ecological*  
50                  *Indicators*. A definitive version was subsequently published as *Ecological Indicators*, Issue 24, 2013,  
51                  DOI 10.1016/j.ecolind.2012.08.005

52 **ABSTRACT**

53 Human demand on ecosystem services continues to increase, and evidence suggests that this demand is  
54 outpacing the regenerative and absorptive capacity of the biosphere. As a result, the productivity of  
55 natural capital may increasingly become a limiting factor for the human endeavor. Metrics tracking  
56 human demand on, and availability of, regenerative and waste absorptive capacity within the biosphere  
57 are therefore needed. Ecological Footprint analysis is such a metric; it measures human appropriation  
58 (Ecological Footprint) and the biosphere's supply (biocapacity) of ecosystem products and services in  
59 terms of the amount of bioproductive land and sea area (ecological assets) needed to supply these  
60 products and services.

61 This paper documents the latest method for estimating the Ecological Footprint and biocapacity of  
62 nations, using the National Footprint Accounts (NFA) applied to more than 200 countries and for the  
63 world overall. Results are also compared with those obtained from previous editions of the NFA.  
64 According to the 2011 Edition of the National Footprint Accounts, humanity demanded the resources  
65 and services of 1.5 planets in 2008; this human demand was 0.7 planets in 1961.

66 Situations in which total demand for ecological goods and services exceed the available supply for a  
67 given location, are called 'overshoot'. 'Global overshoot' indicates that stocks of ecological capital are  
68 depleting and/or that waste is accumulating. As the methodology keeps being improved, each new  
69 edition of the NFA supports the findings of a global overshoot.

70

71 **Keywords:** Ecological Footprint, biocapacity, resource accounting, planetary limits, NFA editions  
72 comparison, sensitivity analysis.

73 **1. Introduction**

74 Economic prosperity and societal well-being depend on the planet's capacity to provide resources and  
75 ecosystem services (e.g., Costanza et al., 1997; Costanza and Daly, 1992; Daly, 1990; Daly and Farley,  
76 2004; DeFries et al., 2004; Max-Neef, 1995). While most policy decisions are made under an  
77 assumption of limitless resources and ecosystem services, the planet has boundaries and sustainable  
78 development cannot be secured without operating within them (Rockström et al., 2009a).

79

80 Environmental changes such as deforestation, collapsing fisheries, and carbon dioxide accumulation in  
81 the atmosphere indicate that human demand is likely to be exceeding the regenerative and absorptive  
82 capacity of the biosphere. As the demands upon natural systems rapidly increase due to the swelling  
83 global economy and the need to attain better standards of living, several studies suggest that many of  
84 the Earth's thresholds are being exceeded and that, because of this, the Biosphere's future ability to  
85 provide for humanity is at risk (Goudie, 1981; Haberl, 2006; Nelson et al., 2006; Moore et al., 2012;  
86 Rockström et al., 2009b; Scheffer et al., 2001; Schlesinger, 2009; Thomas et al., 2004).

87

88 Barnosky et al (2012) argue that a planetary-scale critical transition is approaching as a result of the  
89 many human pressures, and that tools are needed to detect early warning signs and to forecast the  
90 consequences of such pressures on ecosystems. Careful management of human interaction with the  
91 biosphere is thus essential to ensure future prosperity; systemic accounting tools are needed for tracking  
92 the combined effects of the many pressures that humans are placing on the planet (Galli et al., 2012).

93

94 The Ecological Footprint is a potential tool to jointly measure planetary boundaries and the extent to  
95 which humanity is exceeding them. It can be used to investigate issues such as the limits of resource  
96 consumption, the international distribution of the world's natural resources, and how to address the  
97 sustainability of natural resource use across the globe. Assessing current ecological supply and demand  
98 as well as historical trends provides a basis for setting goals, identifying options for action, and tracking  
99 progress toward stated goals.

100

101 The first systematic attempt to calculate the Ecological Footprint and biocapacity of nations began in  
102 1997 (Wackernagel et al. 1997). Building on these assessments, Global Footprint Network initiated its  
103 National Footprint Accounts (NFA) program in 2003, with the most recent Edition issued in 2011.  
104 NFAs constitute an accounting framework quantifying the annual supply of, and demand for, key  
105 ecosystem services by means of two measures (Wackernagel et al., 2002):

106

- 107       • **Ecological Footprint:** a measure of the demand populations and activities place on the  
108       biosphere in a given year, given the prevailing technology and resource management of that  
109       year.
- 110       • **Biocapacity:** a measure of the amount of biologically productive land and sea area available to  
111       provide the ecosystem services that humanity consumes – our ecological budget or nature’s  
112       regenerative capacity.

113

114   Ecological Footprint and biocapacity values are expressed in mutually exclusive units of area necessary  
115   to annually provide (or regenerate) such ecosystem services. They include<sup>1</sup>: cropland for the provision  
116   of plant-based food and fiber products; grazing land and cropland for animal products; fishing grounds  
117   (marine and inland) for fish products; forests for timber and other forest products; uptake land to  
118   neutralize waste emissions (currently only the areas for absorbing anthropogenic carbon dioxide  
119   emissions are considered); and built-up areas for shelter and other infrastructure.

120

121   This paper describes the methodology for calculating the Ecological Footprint and biocapacity utilized  
122   in the 2011 Edition of the National Footprint Accounts and provides researchers and practitioners with  
123   information to deepen their understanding of the calculation methodology. It builds on previous  
124   Ecological Footprint work and methodology papers for the National Footprint Accounts (Wackernagel,  
125   1991; Rees 1992, Wackernagel, 1994; Wackernagel and Rees, 1996; Wackernagel et al. 1997,  
126   Wackernagel et al. 1999a, b, Wackernagel et al. 2002, Monfreda et al. 2004, Wackernagel et al. 2005,  
127   Galli, 2007; Kitzes et al. 2007a, Ewing et al. 2010a). It also compares the most recent Ecological  
128   Footprint and biocapacity results with those from previous editions of the National Footprint Accounts.

129

## 130   **2. National Footprint Accounts: data sources and accounting framework**

131   Global Footprint Network releases National Footprint Accounts (NFA) annually. The NFA 2011  
132   Edition calculate the Ecological Footprint and biocapacity of more than 200 countries and territories, as  
133   well as global totals, from 1961 to 2008 (Global Footprint Network, 2011). The intent of the NFA is to  
134   provide scientifically robust and transparent calculations to highlight the relevance of biocapacity limits  
135   for decision-making. The National Footprint Accounts measure one main aspect of sustainability only -  
136   *how much biocapacity humans demand in comparison to how much is available* - not all aspects of  
137   sustainability, nor all environmental concerns. The attempt to answer this particular scientific research

---

<sup>1</sup> In theory, the Ecological Footprint includes all human demands that compete for space, and biocapacity all areas that provide such services. But in practice, consistent data sets for all aspects do not exist. For this reason not all human demands that compete for space are included in actual assessments, nor all areas that provide services.

138 question is motivated by the assumption that the Earth’s regenerative capacity is the limiting factor for  
 139 the human economy in times when human demand exceeds what the biosphere can renew.

140

141 The calculations in the NFA are based primarily on data sets (Table 1) from UN agencies or affiliated  
 142 organizations such as the Food and Agriculture Organization of the United Nations (FAOSTAT, 2011),  
 143 the UN Statistics Division (UN Commodity Trade Statistics Database – UN Comtrade 2011), and the  
 144 International Energy Agency (IEA 2011). Other data sources include studies in peer-reviewed journals  
 145 and thematic collections.

146

147 TABLE 1: Input data to the Ecological Footprint and biocapacity calculation. Approximately 61 million  
 148 data points are used in the National Footprint Accounts 2011 Edition (6,000 data points per country and  
 149 year).

150

DATASET	SOURCE	DESCRIPTION
Production of primary agricultural products	FAO ProdSTAT	Data on physical quantities (tonnes) of primary products produced in each of the considered countries
Production of crop-based feeds used to feed animals	Feed from general marketed crops data is directly drawn from the SUA/FBS from FAOSTAT Data on crops grown specifically for fodder is drawn directly from the FAO ProdSTAT	Data on physical quantities (tonnes) of feeds, by type of crops, available to feed livestock
Production of seeds	Data on crops used as seeds is calculated by Global Footprint Network based on data from the FAO ProdSTAT	Data on physical quantities (tonnes) of seed
Import and Export of primary and derived agricultural and livestock products	FAO TradeSTAT	Data on physical quantities (tonnes) of products imported and exported by each of the considered countries
Import and Export of non-agricultural commodities	COMTRADE	Data on physical quantities (kg) of products imported and exported by each of the considered countries
Livestock crop consumption	Calculated by Global Footprint Network based upon the following datasets:  <ul style="list-style-type: none"> <li>• FAO Production for primary Livestock</li> <li>• Haberl et al., 2007.</li> </ul>	Data on crop-based feed for livestock (tonnes of dry matter per year), split into different crop categories

Production of primary forestry products as well as import and export of primary and derived forestry products	FAO ForeSTAT	Data on physical quantities (tonnes and m <sup>3</sup> ) of products (timber and wood fuel) produced, imported and exported by each country
Production of primary fishery products as well as import and export of primary and derived fishery products	FAO FishSTAT	Data on physical quantities (tonnes) of marine and inland fish species landed as well as import and export of fish commodities
Carbon dioxide emissions by sector	International Energy Agency (IEA)	Data on total amounts of CO <sub>2</sub> emitted by each sector of a country's economy
Built-up/infrastructure areas	A combination of data sources is used, in the following order of preference:  <ol style="list-style-type: none"> <li>1. CORINE Land Cover</li> <li>2. FAO ResourceSTAT</li> <li>3. Global Agro-Ecological Zones (GAEZ) Model</li> <li>4. Global Land Cover (GLC) 2000</li> <li>5. Global Land Use Database from the Center for Sustainability and the Global Environment (SAGE) at University of Wisconsin</li> </ol>	Built-up areas by infrastructure type and country. Except for data drawn from CORINE for European countries, all other data sources only provide total area values
Cropland yields	FAO ProdSTAT	World average yield for 164 primary crop products
National yield factors for cropland	Calculated by Global Footprint Network based on cropland yields and country specific unharvested percentages	Country specific yield factors for cropland
Grazing land yields	Chad Monfreda (personal communication), 2008. SAGE, University of Wisconsin, Madison	World average yield for grass production. It represents the average above-ground edible net primary production for grassland available for consumption by ruminants
Fish yields	Calculated by Global Footprint Network based on several data sources including: <ul style="list-style-type: none"> <li>• Sustainable catch value (Gulland, 1971)</li> <li>• Trophic levels of fish species (Fishbase Database available at <a href="http://www.fishbase.org">www.fishbase.org</a>)</li> <li>• Data on discard factors, efficiency transfer, and carbon content of fish per tonne wet weight (Pauly and Christensen, 1995)</li> </ul>	World-average yields for fish species. They are based on the annual marine primary production equivalent


Forest yields	World average forest yield calculated by Global Footprint Network based on national Net Annual Increment (NAI) of biomass. NAI data is drawn from two sources: <ul style="list-style-type: none"> <li>• Temperate and Boreal Forest Resource Assessment – TBFRA (UNECE and FAO 2000)</li> <li>• Global Fiber Supply Model – GFSM (FAO, 1998)</li> </ul>	World average forest yield. It is based on the forests' Net Annual Increment of biomass.  NAI is defined as the average annual volume over a given reference period of gross increment less that of neutral losses on all trees to a minimum diameter of 0 cm (d.b.h.)
Carbon Uptake land yield	Calculated by Global Footprint Network based on data on terrestrial carbon sequestration (IPCC 2006) and the ocean sequestration percentage (Khatriwala et al., 2009)  Further details can be found in (Gracey et al., 2012)	World average carbon uptake capacity. Though different ecosystems have the capacity to sequester CO <sub>2</sub> , carbon uptake land is currently assumed to be forest land only by the Ecological Footprint methodology
Equivalence Factors (EQF)	Calculated by Global Footprint Network based on data on land cover and agricultural suitability  Data on agricultural suitability is obtained from the Global Agro-Ecological Zones (GAEZ) model (FAO and IIASA, 2000).  Land cover data drawn from the FAO ResourceSTAT database	EQF for crop, grazing, forest and marine land. Based upon the suitability of land as measured by the Global Agro-Ecological Zones model

151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165

---

Results can be reported at the level of a product category, land use type, or aggregated into a single number (Figure 1) - the latter being the most commonly used reporting format. Normalizing factors, referred as the yield factor and equivalence factor, are used to scale the contribution of each single land use type so that values can be added up into an aggregate number (see sections 4.2 and 4.3). Aggregating results into a single value has the advantage of monitoring the combined demand of anthropogenic activities against nature's overall regenerative capacity. It also helps to understand the complex relationships between the many environmental problems exposing humanity to a "peak-everything" situation (Heinberg, 2007). This is a unique feature since pressures are more commonly evaluated independently (climate change, fisheries collapse, land degradation, land use change, food consumption, etc.).

FIGURE 1: National Footprint Accounts (NFA)’ accounting framework.



167

168 **3. Calculation methodology**

169 *3.1 Ecological Footprint and biocapacity: basic equations*

170 The Ecological Footprint measures appropriated biocapacity across five distinct land use types. This is  
 171 contrasted with six demand categories. The reason for this discrepancy is that two demand categories,  
 172 forest products and carbon sequestration, compete for the same biocapacity category: forest land.  
 173 Average bioproductivity differs between various land use types, as well as between countries for any  
 174 given land use type. For comparability across land use types and countries, Ecological Footprint and  
 175 biocapacity are usually expressed in units of world-average bioproductive area, referred to as global  
 176 hectares (gha).

177 Global hectares provide more information than simply weight - which does not capture the extent of  
 178 land and sea area used - or physical area - which does not capture how much ecological production is  
 179 associated with that land. Two important types of coefficients, the yield factors (YF) and the



180 equivalence factors (EQF), allow results to be expressed in terms of a standardized - cross-country  
 181 comparable - unit of measure (Monfreda et al., 2004; Galli et al., 2007). The use of global hectares  
 182 allows for the addition of Ecological Footprint (and biocapacity) values of different land use types into  
 183 a single number: consumption-focused applications that have a global context, and global sustainability  
 184 studies aiming at comparing the Ecological Footprint (and biocapacity) results of Nations benefit from  
 185 the use of global hectares (Ferguson, 1999; Wackernagel et al., 2004).

186

187 For a given nation, the Ecological Footprint of production,  $EF_P$ , represents primary demand for  
 188 biocapacity and is calculated as

189

$$EF_P = \sum_i \frac{P_i}{Y_{N,i}} \cdot YF_{N,i} \cdot EQF_i = \sum_i \frac{P_i}{Y_{W,i}} \cdot EQF_i \quad (\text{Equation 1})$$

191

192 where  $P$  is the amount of each primary product  $i$  that is harvested (or carbon dioxide emitted) in the  
 193 nation;  $Y_{N,i}$  is the annual national average yield for the production of commodity  $i$  (or its carbon uptake  
 194 capacity in cases where  $P$  is  $CO_2$ );  $YF_{N,i}$  is the country-specific yield factor for the production of each  
 195 product  $i$ ;  $Y_{W,i}$  is the average world yield for commodity  $i$ ; and  $EQF_i$  is the equivalence factor for the  
 196 land use type producing products  $i$ .

197

198 The definition of  $YF_{N,i}$  as the ratio between  $Y_{N,i}$  and  $Y_{W,i}$  (see section 4.2) leads to the equivalence of the  
 199 second and third terms in Equation 1. The latter manifestation of the equation is used in the NFA  
 200 calculations.

201

202 A variety of derived products are also tracked in the NFA (see Table 1), for which production yields  
 203 ( $Y_W$ ) have to be calculated before the implementation of Equation 1. Primary and derived goods are  
 204 related by product specific extraction rates. The extraction rate for a derived product,  $EXTR_D$ , is used to  
 205 calculate its effective yield as follows:

206

$$Y_{W,D} = Y_{W,P} \cdot EXTR_D \quad (\text{Equation 2})$$

208

209 where  $Y_{W,D}$  and  $Y_{W,P}$  are the world-average yield for the derived and the primary product, respectively.

210

211 Often  $EXTR_D$  is simply the mass ratio of derived product to primary input required. This ratio is known  
 212 as the technical conversion factor (FAO, 2000) for the derived product, denoted as  $TCF_D$  below. There  
 213 are a few cases where multiple derived products are created simultaneously from the same primary  
 214 product. For example, soybean oil and soybean cake are both extracted simultaneously from the same  
 215 primary product, in this case soybeans. In this situation, summing the primary product equivalents of  
 216 the derived products would lead to double counting. To resolve this problem, the Ecological Footprint  
 217 of the primary product must be shared between the simultaneously derived goods. The generalized  
 218 formula for the extraction rate for a derived good  $D$  is  
 219

$$EXTR_D = \frac{TCF_D}{FAF_D} \quad \text{(Equation 3)}$$

221 where  $FAF_D$  is the Footprint allocation factor. This allocates the Footprint of a primary product between  
 222 simultaneously derived goods according to the TCF-weighted prices. The prices of derived goods  
 223 represent their relative contributions to the incentive for the harvest of the primary product. This is the  
 224 only point in the entire NFA framework where monetary data is used to allocate physical flows;  
 225 moreover, this method assumes a constant price-to-mass relationship over time, which is unlikely to be  
 226 the case.  
 227

228

229 The equation for the Footprint allocation factor of a derived product is

230

$$FAF_D = \frac{TCF_D \cdot V_D}{\sum TCF_i \cdot V_i} \quad \text{(Equation 4)}$$

232

233 where  $V_i$  is the market price of each simultaneous derived product (2008 market prices were used in the  
 234 NFA 2011 Edition, throughout the whole 1961-2008 period). For a production chain with only one  
 235 derived product, then,  $FAF_D$  is 1 and the extraction rate is equal to the technical conversion factor.  
 236

237

237 For a given country, the biocapacity  $BC$  is calculated as follows:

238

$$BC = \sum_i A_{N,i} \cdot YF_{N,i} \cdot EQF_i \quad \text{(Equation 5)}$$

240

241 where  $A_{N,i}$  is the bioproductive area that is available for the production of each product  $i$  at the country  
 242 level,  $YF_{N,i}$  is the country-specific yield factor for the land producing products  $i$ , and  $EQF_i$  is the  
 243 equivalence factor for the land use type producing each product  $i$ .

244

### 245 3.2 Yield factors

246 Yield factors (YFs) account for countries' differing levels of productivity for particular land use types.<sup>2</sup>  
 247 YFs are country-specific and vary by land use type and year. They may reflect natural factors such as  
 248 differences in precipitation or soil quality, as well as anthropogenic differences such as management  
 249 practices.

250

251 The YF is the ratio of national average to world average yields. It is calculated in terms of the annual  
 252 availability of usable products. For any land use type  $L$ , a country's yield factor  $YF_L$ , is given by

253

$$254 \quad YF_L = \frac{\sum_{i \in U} A_{W,i}}{\sum_{i \in U} A_{N,i}} \quad \text{(Equation 6)}$$

255

256 where  $U$  is the set of all usable primary products that a given land use type yields, and  $A_{W,i}$  and  $A_{N,i}$  are  
 257 the areas necessary to furnish that country's annually available amount of product  $i$  at world and  
 258 national yields, respectively. These areas are calculated as

259

$$260 \quad A_{N,i} = \frac{P_i}{Y_{N,i}} \quad \text{and} \quad A_{W,i} = \frac{P_i}{Y_{W,i}} \quad \text{(Equation 7)}$$

261

262 where  $P_i$  is the total national annual growth of product  $i$ , and  $Y_{N,i}$  and  $Y_{W,i}$  are national and world yields  
 263 for the same product, respectively. Thus  $A_{N,i}$  is always the area that produces a given product  $i$  within a  
 264 given country, while  $A_{W,i}$  gives the equivalent area of world-average land yielding the same product  $i$ .

265

266 With the exception of cropland, all land use types included in the NFAs are assumed to provide only a  
 267 single human-useful primary product  $i$ , such as wood from forest land or grass from grazing land. For  
 268 these land use types, the equation for the YF simplifies to

---

<sup>2</sup> For example, the average hectare of pasture in New Zealand produces more grass than a world average hectare of pasture land. Thus, in terms of productivity, one hectare of grassland in New Zealand is equivalent to more than one world average grazing land hectare; it is potentially capable of supporting more meat production.

269

270

$$YF_L = \frac{Y_{N,i}}{Y_{W,i}}$$

(Equation 8)

271

272 Due to the difficulty of assigning a yield to built-up land, the YF for this land use type is assumed to be  
273 the same as that for cropland; urban areas are assumed to be built on productive agricultural lands. For  
274 lack of detailed global datasets, areas inundated by hydroelectric reservoirs are presumed to have  
275 previously had world average productivity. The YF for the carbon Footprint is assumed to be the same  
276 as that for forest land, due to limited data availability regarding the carbon uptake of other land use  
277 types. All inland waters are assigned a YF of one, due to the lack of a comprehensive global dataset on  
278 freshwater ecosystem productivities.

279

### 280 *3.3 Equivalence factors*

281 In order to combine the Ecological Footprint or biocapacity of different land-use types, a second  
282 coefficient is necessary (Galli et al., 2007). Equivalence factors (EQFs) convert the areas of different  
283 land use types, at their respective world average productivities, into their equivalent areas at global  
284 average bioproductivity across all land use types. EQFs vary by land use type as well as by year.

285

286 The rationale behind the EQF calculation is to weight different land areas in terms of their inherent  
287 capacity to produce human-useful biological resources. The weighting criterion is not the actual  
288 quantity of biomass produced, but what each hectare would be able to inherently deliver.

289 As an approximation of inherent capacity, EQFs are currently calculated<sup>3</sup> using suitability indexes from  
290 the Global Agro-Ecological Zones model combined with data on the actual areas of cropland, forest  
291 land, and grazing land area from FAOSTAT (FAO and IIASA, 2000; FAO ResourceSTAT Statistical  
292 Database 2008). The GAEZ model divides all land globally into five categories, based on calculated  
293 potential crop productivity under assumption of agricultural input. All land is assigned a quantitative  
294 suitability index from among the following:

295

- 296 • Very Suitable (VS) – 0.9

---

<sup>3</sup> Actual Net Primary Production (NPP) values have been suggested for use in scaling land type productivity (Venetoulis and Talberth, 2008) and were also used in the earliest Footprint accounts; however, this would not allow incorporating the inherent productivity as, for instance, crop land is managed for maximum crop, not for maximum biomass production. Potential NPP data - the NPP of useable biological materials that could be potentially available in the absence of human management - could theoretically be used as weighting factors (see Kitzes et al., 2009). A global data set exists (FAO, 2006) and research is under way at Global Footprint Network to assess the possibility of using potential NPP data in calculating EQFs.

- 297 • Suitable (S) – 0.7
- 298 • Moderately Suitable (MS) – 0.5
- 299 • Marginally Suitable (mS) – 0.3
- 300 • Not Suitable (NS) – 0.1


301

302 The calculation of the EQFs assumes that within each country, the most suitable land available will be  
 303 planted to cropland, after which the most suitable remaining land will be under forest land, and the least  
 304 suitable land will be devoted to grazing land (Wackernagel et al., 2002). In each year, EQFs are  
 305 calculated as the ratio of the world average suitability index for a given land use type to the average  
 306 suitability index for all land use types. Figure 2 shows a schematic of this calculation.

307

308 **FIGURE 2: Schematic Representation of equivalence factor calculations.**

309



310

311

312 The total number of bioproductive land hectares is shown by the length of the horizontal axis. Vertical  
 313 dashed lines divide this total land area into the three terrestrial land use types for which equivalence  
 314 factors are calculated (cropland, forest, and grazing land). The length of each horizontal bar in the graph  
 315 shows the total amount of land available with each suitability index. The vertical location of each bar  
 316 reflects the suitability score for that suitability index, between 10 and 90.

317

318 For the reasons detailed above, the EQF for built-up land is set equal to that for cropland, except when  
319 there is clear evidence that built-up land does not sit on cropland. EQF of carbon uptake land is set  
320 equal to that of forest land since the carbon Footprint is assumed to draw on forest area. The EQF for  
321 hydroelectric reservoir area is set equal to one, reflecting the assumption that hydroelectric reservoirs  
322 flood world average bioproductive land. The EQF for marine area is calculated such that the amount of  
323 calories of salmon that can be produced by a single global hectare of marine area will be equal to the  
324 amount of calories of beef produced by a single global hectare of pasture. This is based on the  
325 assumption that a calorie from animal protein from land and from sea would be considered to be of  
326 equivalent resource value for human consumption. The EQF for inland water is set equal to that of  
327 marine area.

328

### 329 *3.4 A Consumer approach for the National Footprint Accounts*

330 All manufacturing processes rely to some degree on the use of biocapacity to provide material inputs  
331 and remove wastes at various points in the production chain. Thus all products carry with them an  
332 embodied Footprint and international trade flows can be seen as flows of embodied demand for  
333 biocapacity (see Figure 3).

334

335 In order to keep track of the biocapacity - both direct and indirect - needed to support people's  
336 consumption patterns and to properly allocate the Footprints of traded goods to final consumers, the  
337 National Footprint Accounts use a consumer-based approach; for each land use type, the Ecological  
338 Footprint of consumption ( $EF_C$ ) is thus calculated as

339

$$340 \quad EF_C = EF_P + EF_I - EF_E \quad \text{(Equation 9)}$$

341

342 where  $EF_P$  is the Ecological Footprint of production and  $EF_I$  and  $EF_E$  are the Footprints embodied in  
343 imported and exported commodity flows, respectively. For each traded product,  $EF_I$  and  $EF_E$  are  
344 calculated as in equation 1, with production  $P$  being the amount of product  $i$  imported or exported,  
345 respectively.

346

347

348

349

350


351

352

353

354

FIGURE 3: Schematic of direct and indirect demand for domestic and global biocapacity.



355

356


#### 357 4. Land use types in the National Footprint Accounts

358 The Ecological Footprint represents demand for ecosystem products and services in terms of  
 359 appropriation of various land use types (see Section 1) while biocapacity represents the productivity  
 360 available to serve each use. In 2008, the area of biologically productive land and water on Earth was  
 361 approximately 12 billion hectares. After multiplying by the EQFs, the relative area of each land use  
 362 type expressed in global hectares differs from the distribution in actual hectares as shown in Figure 4.

363

364 FIGURE 4: Relative area of land use types worldwide in hectares and global hectares, 2008.

365



366

367

368 4.1 Cropland

369 Cropland<sup>4</sup> consists of the area required to grow all crop products, including livestock feeds, fish meals,  
370 oil crops and rubber. It is the most bioproductive of the land use types included in the NFAs. In other  
371 words, the number of global hectares of cropland is large compared to the number of physical hectares  
372 of cropland in the world, as shown in Figure 4.

373

374 NFAs calculate the Footprint of cropland using data on production, import and export of primary and  
375 derived agricultural products. The Footprint of each crop type is calculated as the area of cropland that  
376 would be required to produce the harvested quantity at world-average yields.

377

378 Cropland biocapacity represents the combined productivity of all land devoted to growing crops, which  
379 the cropland Footprint cannot exceed. As an actively managed land use type, cropland has yields of  
380 harvest equal to yields of growth by definition and thus it is not possible for the Footprint of production  
381 of this land use type to exceed biocapacity within any given area (Kitzes et al., 2009). The eventual  
382 availability of data on present and historical sustainable crop yields would allow for improving the  
383 cropland footprint calculation and tracking crop overexploitation (Bastianoni et al., 2012).

384

385 4.2 Grazing Land

386 The grazing land Footprint measures the area of grassland used in addition to crop feeds to support  
387 livestock. Grazing land<sup>5</sup> comprises all grasslands used to provide feed for animals, including cultivated  
388 pastures as well as wild grasslands and prairies. The grazing land Footprint is calculated following  
389 Equation 1, where yield represents average above-ground NPP for grassland. The total demand for  
390 pasture grass,  $P_{GR}$ , is the amount of biomass required by livestock after cropped feeds are accounted for,  
391 following the formula

392

$$P_{GR} = TFR - F_{Mkt} - F_{Crop} - F_{Res} \quad (\text{Equation 10})$$

394

395 where  $TFR$  is the calculated total feed requirement, and  $F_{Mkt}$ ,  $F_{Crop}$  and  $F_{Res}$  are the amounts of feed  
396 available from general marketed crops, crops grown specifically for fodder, and crop residues,  
397 respectively.

---

<sup>4</sup> In the National Footprint Accounts, “cropland” is defined to match the FAO land use category ‘Arable land and Permanent crops’ – FAO code 6620.

<sup>5</sup> In the National Footprint Accounts, “grazing land” is defined to match the FAO land use category ‘Permanent meadows and pastures’ – FAO code 6655.



398

399 The grazing land calculation is the most complex in the NFAs and significant improvements have taken  
400 place over the past seven years; including improvements to the total feed requirement, the inclusion of  
401 fish and animal products used as livestock feed, and the inclusion of livestock food aid (see Ewing et  
402 al., 2010a for further details).

403

404 However, as the yield of grazing land represents the amount of above-ground primary production  
405 available in a year with no significant prior stocks to draw down, and given the fact that soil depletion is  
406 not tracked by the Ecological Footprint methodology (Kitzes et al., 2009), an eventual overshoot for  
407 this land use type still cannot be shown.

408

#### 409 4.3 Fishing Grounds

410 The fishing grounds Footprint is calculated based on the annual primary production required to sustain a  
411 harvested aquatic species. This primary production requirement, denoted *PPR*, is the mass ratio of  
412 harvested fish to annual primary production needed to sustain that species, based on its average trophic  
413 level. Equation 11 provides the formula used to calculate *PPR*. It is based on the work of Pauly and  
414 Christensen (1995).

415

$$416 \quad PPR = CC \cdot DR \cdot \left( \frac{1}{TE} \right)^{(TL-1)} \quad \text{(Equation 11)}$$

417

418 where *CC* is the carbon content of wet-weight fish biomass, *DR* is the discard rate for bycatch, *TE* is the  
419 transfer efficiency of biomass between trophic levels, and *TL* is the trophic level of the fish species in  
420 question.

421

422 In the NFAs, *DR* is assigned the global average value of 1.27 for all fish species, meaning that for every  
423 ton of fish harvested, 0.27 tonnes of bycatch are also harvested (Pauly and Christensen 1995). This  
424 bycatch rate is applied as a constant coefficient in the *PPR* equation, reflecting the assumption that the  
425 trophic level of bycatch is the same as that of the primary catch species. These approximations are  
426 employed for lack of higher resolution data on bycatch. *TE* is assumed to be 0.1 for all fish, meaning  
427 that 10% of biomass is transferred between successive trophic levels (Pauly and Christensen, 1995).

428

429 The estimate of annually available primary production used to calculate marine yields is based on  
430 estimates of the sustainable annual harvests of 19 different aquatic species groups (Gulland, 1971).

431 These quantities are converted to primary production equivalents using Equation 11, and the sum of  
432 these is taken to be the total primary production requirement that global fisheries may sustainably  
433 harvest. Thus the total sustainably harvestable primary production requirement,  $PP_S$ , is calculated as  
434

$$435 \quad PP_S = \sum(Q_{S,i} \cdot PPR_i) \quad \text{(Equation 12)}$$

436  
437 where  $Q_{S,i}$  is the estimated sustainable catch for species group  $i$ , and  $PPR_i$  is the primary production  
438 requirement corresponding to the average trophic level of species group  $i$ . Thus the world-average  
439 marine yield  $Y_M$ , in terms of PPR, is given by  
440

$$441 \quad Y_M = \frac{PP_S}{A_{CS}} \quad \text{(Equation 13)}$$

442  
443 where  $PP_S$  is the global sustainable harvest from Equation 12, and  $A_{CS}$  is the global total continental  
444 shelf area.

445  
446 Significant improvements have taken place over the past seven years in the calculation of the fishing  
447 grounds section of the NFAs, including the revision of many fish extraction rates, the inclusion of  
448 aquaculture production and of crops used in aquafeeds (see Ewing et al., 2010a for further).

#### 449 450 *4.4 Forest Land*

451 The forest land<sup>6</sup> Footprint measures the annual harvest of fuel wood and timber to supply forest  
452 products. The yield used in the forest land Footprint is the net annual increment (NAI) of merchantable  
453 timber per hectare. Timber productivity data from the UNEC and FAO Forest Resource Assessment and  
454 the FAO Global Fiber Supply are utilized to calculate the world average yield of 1.81 m<sup>3</sup> of harvestable  
455 wood per hectare per year (UNECE and FAO 2000; FAO 1998).

456  
457 NFAs calculate the Footprint of forest land according to the production quantities of 13 primary timber  
458 products and three wood fuel products. Trade flows include 30 timber products and 3 wood fuel  
459 products.

460

---

<sup>6</sup> In the National Footprint Accounts, “forest” is defined to match the FAO land use category ‘Forest Area’ – FAO code 6661. Due to data limitation, current accounts do not distinguish between forests for forest products, for long-term carbon uptake, or for biodiversity reserves.

461 4.5 Carbon Footprint

462 The uptake land to accommodate the carbon Footprint is the only land use type included in the NFAs  
463 that is exclusively dedicated to tracking a waste product: carbon dioxide.<sup>7</sup> In addition, it is the only land  
464 use type for which biocapacity is currently not explicitly defined. Many different ecosystem types have  
465 the capacity for long-term storage of CO<sub>2</sub>, including land use types such as cropland or grassland.  
466 However, since most terrestrial carbon uptake in the biosphere occurs in forests, and to avoid  
467 overestimations, carbon uptake land is assumed to be (a subcategory of) forest land by the Ecological  
468 Footprint methodology. Therefore, forest for timber and fuelwood is not separated from forest for  
469 carbon uptake.<sup>8</sup>

470

471 CO<sub>2</sub> is released into the atmosphere from a variety of sources, including human activities such as  
472 burning fossil fuels and certain land use practices; as well as natural events such as forest fires,  
473 volcanoes, and respiration by animals and microbes. Analogous to Equation 1, the formula for the  
474 carbon Ecological Footprint ( $EF_C$ ) is

475

$EF_C = \frac{P_C \cdot (1 - S_{Ocean})}{Y_C} * EQF$	(Equation 14)
--	---------------

477

478 where  $P_C$  is the annual anthropogenic emissions (production) of carbon dioxide,  $S_{Ocean}$  is the fraction of  
479 anthropogenic emissions sequestered by oceans in a given year (see section 5.3 for further details) and  
480  $Y_C$  is the annual rate of carbon uptake per hectare of world average forest land.

481

482 4.6 Built-Up Land

483 The built-up land Footprint is calculated based on the area of land covered by human infrastructure:  
484 transportation, housing, industrial structures and reservoirs for hydroelectric power generation. The  
485 NFA 2011 Edition assumes that built-up land occupies what would previously have been cropland,  
486 except in cases where evidences exist that built-up land does not sit on cropland (e.g., in the United  
487 Arab Emirates – see Abdullatif and Alam, 2011). This assumption is based on the observation that

---

<sup>7</sup> Today, the term “carbon footprint” is widely used as shorthand for the amount of anthropogenic greenhouse gas emissions; in the Ecological Footprint methodology however, it translates the amount of anthropogenic carbon dioxide into the amount of productive land and sea area required to sequester carbon dioxide emissions. (See Galli et al. (2012) for additional information.)

<sup>8</sup> Human demands for forest products and carbon uptake capacity are competing for forest land. However, when a forest is used for products, CO<sub>2</sub> is released again in the atmosphere; as such, only legally protected forests with a commitment to long term storage of carbon can truly be counted as uptake areas. Global Footprint Network has not yet identified reliable global data sets on how much of the forest areas are legally protected and dedicated to long-term carbon uptake. For this reason, current National Footprint Accounts do not include a carbon uptake category within the biocapacity calculation.

488 human settlements are generally situated in fertile areas with the potential for supporting high yielding  
489 cropland (Imhoff et al., 1997; Wackernagel et al., 2002).

490 For lack of a comprehensive global dataset on hydroelectric reservoirs, NFAs assume these to cover  
491 world-average bioproductive areas in proportion to their rated generating capacity. Built-up land always  
492 has a biocapacity equal to its Footprint since both quantities capture the amount of bioproductivity lost  
493 to encroachment by physical infrastructure. In addition, the Footprint of production and the Footprint of  
494 consumption of built-up land are always equal in the NFAs as built-up land embodied in traded goods is  
495 not currently included in the calculation due to lack of data. This omission is likely to cause  
496 overestimates of the built-up Footprint of net exporting countries and underestimates of the built-up  
497 Footprint of net importing countries.

498

## 499 **5. Methodological changes between the 2010 and 2011 edition of the National Footprint Accounts**

500 A formal process is in place to assure continuous improvement of the National Footprint Accounts  
501 (NFA) methodology. Coordinated by Global Footprint Network, this process is supported by its  
502 partners and by the National Footprint Accounts Review Committee, as well as other stakeholders.

503

504 There have been three primary motivations for revisions to the NFAs calculation method:

- 505 • to adapt to changes in the organization of the source data;
- 506 • to respond to issues raised in outside reviews; and
- 507 • to increase the detail and accuracy of the NFA calculations.

508

509 This section describes each of the method changes implemented since the NFA 2010 Edition.

510

### 511 *5.1 Data Cleaning*

512 In the NFA 2011, a source data cleaning algorithm was implemented different to the algorithm used in  
513 NFA 2010. The new algorithm is used to reduce (1) spikes and troughs and (2) inconsistencies in the  
514 reported time series of source data sets. The new algorithm excludes data points that are beyond a fixed  
515 distance from the median value of the reference time series data. The fixed distance is a user-defined  
516 multiple of the median value of the time series in question. To replace the removed outliers and/or to  
517 fill in data gaps for non-endpoints, the algorithm interpolates the average value of the two neighbouring  
518 points. To replace endpoints (outliers or missing data), the algorithm extrapolates values based on the  
519 Akaike Information Criterion (Akaike, 1978). The data cleaning algorithm was implemented on the

520 following trade datasets used in the NFA 2011 Edition: the COMTRADE dataset, the FishSTAT  
521 Commodity dataset, and the TRADESTAT dataset from FAOSTAT.

522

523 *5.2 Constant global hectares: a revised method to calculate Ecological Footprint and biocapacity time*  
524 *series*

525 Ecological Footprint and biocapacity calculations are usually presented in units of global hectares (see  
526 section 4). Historically, Ecological Footprint analyses have utilized a Yield Factor (YF) for each land  
527 use type to capture the difference between local and global productivity. The various land use types are  
528 then converted into global hectares using equivalence factors (EQFs) for each land use type. In every  
529 year, the total biocapacity of the planet, expressed in global hectares, equals the total number of  
530 biologically productive physical hectares on Earth (Kitzes et al., 2007b). Therefore, the number of  
531 global hectares of biocapacity available on the planet in any given year only reflects the total physical  
532 bioproductive area of the planet and is entirely insensitive to changes in yields (Wackernagel et al.,  
533 2004). This can cause difficulties of interpretation when comparing changes in biocapacity and  
534 Ecological Footprint over time as it is hard to represent actual variations in demand and supply of  
535 regenerative capacity (Haberl et al., 2001).

536

537 In the NFA 2011 Edition, we have implemented a method for reporting Ecological Footprint and  
538 biocapacity time trends in ‘constant global hectares’ (hectares normalized to have world-average  
539 bioproductivity in a single reference year). This is realized via a set of world-average Intertemporal  
540 Yield Factors (IYFs). By expressing results through the constant global hectare approach, it is possible  
541 to clearly distinguish trends in both total bioproductive area and trends in yield and productivity. IYFs  
542 are calculated for each year and land use type in order to track changes in the world-average  
543 bioproductivity over time of each land type.

544

545 For any given land type producing products  $i$ , in a given year  $j$ , with a selected base year  $b$ , a world  
546 average Intertemporal Yield Factor ( $IYF_w$ ) is thus calculated as:

547

$$IYF_{w,j} = \frac{\sum_i \frac{P_{w,i,j}}{Y_{w,i,b}}}{\sum_i \frac{P_{w,i,j}}{Y_{w,i,j}}} \quad \text{(Equation 15)}$$

549

550 where  $P$  is the amount of a product harvested (or CO<sub>2</sub> emitted) and  $Y_W$  is the world-average product-  
 551 specific yield. For the NFA 2011 Edition, the selected base year is 2008 (the most recent year over the  
 552 analyzed period).

553

554 IYFs complement the function of the Yield Factors (YF) currently employed in the NFAs. While YFs  
 555 compare the yield of a given land use type in a given nation with the world-average yield for that same  
 556 land use type, IYFs account for changes in the world-average yield of that same land use type over  
 557 time.

558

559 Ecological Footprint time series are therefore calculated as follows:

560

$$561 \quad EF = \sum_i \frac{P_{N,i,j}}{Y_{N,i,j}} \cdot YF_{N,i,j} \cdot IYF_{W,i,j} \cdot EQF_{i,j} = \sum_i \frac{P_{N,i,j}}{Y_{W,i,j}} \cdot IYF_{W,i,j} \cdot EQF_{i,j} \quad (\text{Equation 16})$$

562

563 Similarly, biocapacity time series are calculated in terms of constant gha as follows:

564

$$565 \quad BC = \sum_i A_{N,i,j} \cdot YF_{N,i,j} \cdot IYF_{W,i,j} \cdot EQF_{i,j} \quad (\text{Equation 17})$$

566

567 Where, for any product  $i$ , in a given year  $j$ ,  $A_N$  represents the bioproductive area available at the country  
 568 level, and  $YF_N$ ,  $IYF_W$  and  $EQF$ , are the country-specific yield factor, the world average Intertemporal  
 569 Yield Factor, and the equivalence factor for the land use type producing that product, respectively.

570

571 Calculating IYFs for each land use type requires production quantity and yield data over time. While  
 572 production quantity data is available for all products tracked by the NFAs over the period 1961-2008,  
 573 time series yield data are available for crop-based products only. This renders the calculation of IYFs  
 574 currently possible for the ‘cropland’ land use type only; in the absence of available data, IYF time series  
 575 values for all other land types have been set equal to 1.

576

### 577 *5.3 Ocean Uptake Changes*

578 A fraction of human-induced carbon emissions is annually taken up by the oceans from the atmosphere.  
 579 To track this fraction, recent editions of the NFAs have used an averaged ocean uptake value of 1.8 Pg  
 580 C yr<sup>-1</sup> based on two data points drawn from the third IPCC assessment report (IPCC, 2001). This  
 581 quantity has been held constant over time leading to an estimated 82% of anthropogenic emissions

582 taken up by the ocean in 1961, which is likely to be unrealistic. This caused an underestimation of the  
583 carbon Footprint component in the early decades tracked by the NFAs.

584 To create an appropriate time series for the percent uptake of anthropogenic carbon emissions into the  
585 ocean, in the NFA 2011 Edition we have used ocean uptake data (in Pg C yr<sup>-1</sup>) from Khatiwala et al  
586 (2009) and divided this data by the corresponding (total anthropogenic) carbon emissions data (in Pg C  
587 yr<sup>-1</sup>) from the Carbon Dioxide Information Analysis Center (Marland et al., 2007). The outcome of the  
588 revised calculation shows a relatively constant percentage uptake for oceans, varying between 28% and  
589 35% over the period 1961-2008.

590 Implementing this change has caused a major shift in the total humanity's Footprint value from 1961 to  
591 the late 1990s; this has significantly contributed to a shift in the global overshoot state - the first  
592 occurrence of overshoot is calculated as occurring in the early 1970s (NFA 2011 Edition), changed  
593 from the mid 1970s (NFA 2010 Edition).

594

## 595 **6. Results**

596 According to the 2011 Edition of the National Footprint Accounts, in 1961 humanity's Ecological  
597 Footprint was approximately half of what the biosphere could supply annually; humanity was living off  
598 the planet's annual ecological interest, not drawing down its principal (Figure 5). Since then,  
599 humanity's overall Footprint has more than doubled, first exceeding the planet's biocapacity in the early  
600 1970s. This situation, known as overshoot, has continued to increase, reaching 52% in 2008.

601

602 In 2008, humanity's Ecological Footprint consisted of 22% cropland, 8% grazing land, 10% forest land,  
603 4% fishing ground, 54% carbon uptake land, and 2% built-up land. As these annual "biocapacity  
604 deficits" accrue into an ever larger ecological debt, ecological reserves are depleting, and wastes such as  
605 CO<sub>2</sub> are accumulating in the biosphere and atmosphere.

606

607

608

609

610

611

612


613

614

615

616

617 FIGURE 5: World overshoot according to the 2011 Edition of the National Footprint Accounts.  
 618 Humanity's Ecological Footprint, expressed in number of planets demanded, has increased significantly  
 619 over the past 47 years.  
 620



621  
 622  
 623  
 624  
 625  
 626  
 627  
 628

Per capita Ecological Footprint and biocapacity results for all countries for the past two years are reported in Table 2 and 3. These tables contain an ordinal ranking of countries by Footprint and biocapacity respectively, as well as a comparison with values from the previous NFA 2010 Edition.

TABLE 2: NFA 2010 and NFA 2011 Editions comparison: Ecological Footprint data table.

	(a) Ecological Footprint 2008 (2011 Edition) [gha/capita]	(b) Ecological Footprint 2007 (2010 Edition) [gha/capita]	(c) Ecological Footprint 2007 (2011 Edition) [gha/capita]	(c)-(b) (c) %	(a)- (c) (a) %	Rank (a)	Rank (b)	Rank (c)	$\Delta$ Rank between editions - (b) to (c)	$\Delta$ Rank between years - (c) to (a)
Afghanistan	0.54	0.62	0.54	-15.30	0.00	156	156	157	-1	1
Albania	1.81	1.91	2.13	10.35	-0.18	89	84	78	6	-11
Algeria	1.65	1.59	1.55	-2.44	0.06	99	101	101	0	2
Angola	0.89	1.00	0.86	-16.92	0.04	140	138	146	-8	6
Argentina	2.71	2.60	2.60	0.19	0.04	66	71	69	2	3
Armenia	1.73	1.75	1.60	-9.32	0.08	94	95	100	-5	6
Australia	6.68	6.84	5.89	-16.11	0.12	8	10	14	-4	6
Austria	5.29	5.30	5.23	-1.30	0.01	19	25	24	1	5
Azerbaijan	1.97	1.87	1.87	0.11	0.05	82	87	82	5	0
Bahrain	6.65	10.04	7.58	-32.41	-0.14	9	3	5	-2	-4
Bangladesh	0.66	0.62	0.66	6.54	-0.01	153	157	155	2	2
Belarus	3.99	3.80	3.70	-2.83	0.07	44	48	47	1	3
Belgium	7.11	8.00	7.39	-8.25	-0.04	7	5	7	-2	0
Benin	1.36	1.23	1.34	8.21	0.01	116	124	113	11	-3
Bolivia	2.61	2.57	2.56	-0.40	0.02	69	72	70	2	1
Bosnia/Herzegovina	2.74	2.75	2.63	-4.70	0.04	65	64	67	-3	2
Botswana	2.84	2.68	2.44	-9.61	0.14	63	69	72	-3	9



Brazil	2.93	2.91	2.80	-3.65	0.04	60	58	57	1	-3
Bulgaria	3.56	4.07	3.02	-34.95	0.15	51	45	53	-8	2
Burkina Faso	1.53	1.32	1.31	-0.45	0.14	104	117	118	-1	14
Burundi	0.85	0.90	0.90	-0.34	-0.07	144	148	143	5	-1
Cambodia	1.19	1.03	1.09	5.51	0.08	123	135	130	5	7
Cameroon	1.09	1.04	1.10	4.90	-0.01	132	134	129	5	-3
Canada	6.43	7.01	6.33	-10.76	0.01	10	8	8	0	-2
C. African Republic	1.36	1.32	1.37	3.53	-0.01	115	116	109	7	-6
Chad	1.89	1.73	1.85	6.97	0.02	85	98	83	15	-2
Chile	3.24	3.24	3.27	1.08	-0.01	53	52	50	2	-3
China	2.13	2.21	2.03	-9.13	0.05	77	76	81	-5	4
Colombia	1.80	1.87	1.80	-3.63	0.00	90	88	88	0	-2
Congo	1.08	0.96	1.01	4.85	0.06	133	141	136	5	3
Costa Rica	2.52	2.69	2.61	-2.89	-0.04	73	67	68	-1	-5
Côte d'Ivoire	0.99	1.01	1.04	3.08	-0.05	136	137	135	2	-1
Croatia	4.19	3.75	3.80	1.36	0.09	39	49	46	3	7
Cuba	1.90	1.85	1.74	-6.57	0.08	84	89	90	-1	6
Cyprus	4.44	6.87	4.37	-57.28	0.02	35	9	39	-30	4
Czech Republic	5.27	5.73	5.38	-6.47	-0.02	20	16	19	-3	-1
Korea, DPR	1.31	1.32	1.31	-0.80	-0.01	118	115	117	-2	-1
Congo, DR	0.76	0.75	0.76	1.47	-0.01	150	152	151	1	1
Denmark	8.25	8.26	8.48	2.57	-0.03	4	4	3	1	-1
Dominican Rep.	1.42	1.47	1.42	-4.05	0.00	110	107	106	1	-4
Ecuador	2.37	1.89	2.11	10.51	0.11	75	86	79	7	4
Egypt	1.70	1.66	1.71	3.21	-0.01	98	100	93	7	-5
El Salvador	1.99	2.03	2.06	1.33	-0.03	81	80	80	0	-1
Eritrea	0.66	0.89	0.91	3.03	-0.39	154	149	142	7	-12
Estonia	4.73	7.88	5.78	-36.29	-0.22	28	7	15	-8	-13
Ethiopia	1.13	1.10	1.08	-1.80	0.04	128	130	133	-3	5
Finland	6.21	6.16	5.96	-3.30	0.04	13	14	13	1	0
France	4.91	5.01	4.86	-3.17	0.01	25	30	29	1	4
Gabon	1.81	1.41	1.68	16.05	0.07	88	110	95	15	7
Gambia	1.41	3.45	1.38	-149.06	0.02	112	51	107	-56	-5
Georgia	1.43	1.82	1.51	-20.31	-0.06	109	90	103	-13	-6
Germany	4.57	5.08	4.69	-8.32	-0.03	32	28	33	-5	1
Ghana	1.74	1.75	1.66	-5.34	0.04	93	94	97	-3	4
Greece	4.92	5.39	5.12	-5.27	-0.04	24	22	27	-5	3
Guatemala	1.78	1.77	1.84	3.35	-0.03	91	93	86	7	-5
Guinea	1.72	1.67	1.67	0.43	0.02	97	99	96	3	-1
Guinea-Bissau	1.10	0.96	1.08	11.24	0.02	131	142	132	10	1
Haiti	0.60	0.68	0.63	-7.00	-0.06	155	155	156	-1	1
Honduras	1.73	1.91	1.83	-4.26	-0.06	95	83	87	-4	-8
Hungary	3.59	2.99	2.99	-0.08	0.17	50	57	54	3	4
India	0.87	0.91	0.86	-6.07	0.01	143	145	145	0	2
Indonesia	1.13	1.21	1.11	-9.49	0.02	130	127	128	-1	-2
Iran, Islamic Rep.	2.66	2.68	2.70	0.50	-0.01	68	68	62	6	-6
Iraq	1.42	1.35	1.45	7.04	-0.02	111	114	104	10	-7
Ireland	6.22	6.29	6.10	-3.15	0.02	12	12	12	0	0
Israel	3.96	4.82	4.28	-12.55	-0.08	46	37	41	-4	-5
Italy	4.52	4.99	4.70	-6.18	-0.04	34	31	32	-1	-2
Jamaica	1.72	1.93	2.22	13.34	-0.29	96	82	77	5	-19
Japan	4.17	4.73	4.37	-8.12	-0.05	40	38	38	0	-2
Jordan	2.13	2.05	2.22	7.71	-0.04	76	79	76	3	0
Kazakhstan	4.14	4.54	4.38	-3.83	-0.06	41	40	37	3	-4
Kenya	0.95	1.11	1.06	-5.01	-0.12	138	129	134	-5	-4
Kuwait	9.72	6.32	7.54	16.07	0.22	2	11	6	5	4
Kyrgyzstan	1.29	1.25	1.22	-2.01	0.05	120	123	122	1	2
Laos	1.30	1.28	1.26	-1.48	0.03	119	119	119	0	0
Latvia	3.95	5.64	5.39	-4.65	-0.36	47	18	18	0	-29
Lebanon	2.85	2.90	2.64	-9.77	0.07	62	59	66	-7	4
Lesotho	1.07	1.07	1.12	4.02	-0.04	134	132	127	5	-7
Liberia	1.28	1.26	1.32	4.30	-0.03	121	120	114	6	-7
Libya	3.19	3.05	3.04	-0.51	0.05	55	55	52	3	-3
Lithuania	4.38	4.67	4.43	-5.47	-0.01	37	39	35	4	-2
Madagascar	1.16	1.79	1.16	-54.60	0.00	127	91	126	-35	-1
Malawi	0.78	0.73	0.78	6.28	-0.01	148	154	149	5	1
Malaysia	3.90	4.86	3.44	-41.42	0.12	49	36	48	-12	-1
Mali	1.86	1.93	1.72	-12.18	0.08	86	81	92	-11	6

Mauritania	2.86	2.61	2.74	4.67	0.04	61	70	58	12	-3
Mauritius	4.55	4.26	4.39	2.90	0.04	33	44	36	8	3
Mexico	3.30	3.00	2.92	-2.68	0.12	52	56	55	1	3
Mongolia	5.53	5.53	5.38	-2.80	0.03	17	20	20	0	3
Morocco	1.32	1.22	1.19	-2.63	0.10	117	125	125	0	8
Mozambique	0.78	0.77	0.76	-1.34	0.03	147	150	153	-3	6
Myanmar	1.94	1.79	1.78	-0.38	0.08	83	92	89	3	6
Namibia	2.03	2.15	2.41	10.64	-0.19	79	77	73	4	-6
Nepal	0.76	3.56	0.76	-367.06	0.00	149	50	152	-102	3
Netherlands	6.34	6.19	6.24	0.72	0.02	11	13	10	3	-1
New Zealand	4.31	4.89	4.20	-16.36	0.03	38	34	42	-8	4
Nicaragua	1.56	1.56	1.63	4.35	-0.04	102	102	98	4	-4
Nigeria	1.44	1.44	1.54	6.97	-0.07	108	108	102	6	-6
Norway	4.77	5.56	5.25	-5.94	-0.10	26	19	23	-4	-3
Palestinian Terr.	0.46	0.74	0.69	-7.02	-0.50	158	153	154	-1	-4
Oman	5.69	4.99	5.36	7.04	0.06	16	32	21	11	5
Pakistan	0.75	0.77	0.77	0.66	-0.02	151	151	150	1	-1
Panama	2.97	2.87	2.87	-0.26	0.03	59	63	56	7	-3
Papua New Guinea	2.68	2.14	2.65	19.35	0.01	67	78	64	14	-3
Paraguay	2.99	3.19	3.28	2.70	-0.10	58	53	49	4	-9
Peru	2.03	1.54	1.85	17.06	0.09	80	103	84	19	4
Philippines	0.98	1.30	0.98	-32.05	0.00	137	118	137	-19	0
Poland	3.94	4.35	4.00	-8.61	-0.02	48	43	44	-1	-4
Portugal	4.12	4.47	4.32	-3.35	-0.05	42	41	40	1	-2
Puerto Rico	0.03	0.04	0.03	-10.34	-0.20	159	159	159	0	0
Qatar	11.68	10.51	10.52	0.13	0.10	1	2	1	1	0
Korea, Rep.	4.62	4.87	4.65	-4.80	-0.01	31	35	34	1	3
Moldova	2.10	1.39	1.37	-1.44	0.35	78	113	108	5	30
Romania	2.84	2.71	2.49	-8.60	0.12	64	65	71	-6	7
Russia	4.40	4.41	4.19	-5.33	0.05	36	42	43	-1	7
Rwanda	0.71	1.02	0.98	-4.41	-0.38	152	136	139	-3	-13
Saudi Arabia	3.99	5.13	4.77	-7.61	-0.20	43	26	31	-5	-12
Senegal	1.53	1.09	1.22	10.02	0.21	103	131	124	7	21
Serbia	2.57	2.39	2.34	-1.89	0.09	71	73	74	-1	3
Sierra Leone	1.13	1.05	1.09	3.49	0.04	129	133	131	2	2
Singapore	6.10	5.34	5.70	6.45	0.07	14	23	16	7	2
Slovakia	4.66	4.06	3.11	-30.58	0.33	30	46	51	-5	21
Slovenia	5.21	5.30	5.15	-2.97	0.01	21	24	26	-2	5
Somalia	1.44	1.42	1.45	1.50	0.00	107	109	105	4	-2
South Africa	2.59	2.32	2.70	13.98	-0.04	70	75	63	12	-7
Spain	4.74	5.42	5.09	-6.58	-0.07	27	21	28	-7	1
Sri Lanka	1.21	1.21	1.22	0.59	-0.01	122	126	123	3	1
Sudan	1.63	1.73	1.72	-0.52	-0.06	100	97	91	6	-9
Swaziland	1.45	1.50	1.24	-20.35	0.14	106	106	120	-14	14
Sweden	5.71	5.88	6.28	6.38	-0.10	15	15	9	6	-6
Switzerland	5.01	5.02	5.28	5.06	-0.05	23	29	22	7	-1
Syria	1.45	1.52	1.35	-12.76	0.07	105	105	112	-7	7
Tajikistan	0.90	1.00	0.87	-14.63	0.03	139	139	144	-5	5
Thailand	2.41	2.37	2.29	-3.34	0.05	74	74	75	-1	1
Macedonia TFYR	5.36	5.66	5.54	-2.12	-0.03	18	17	17	0	-1
Timor-Leste	0.47	0.44	0.53	18.04	-0.13	157	158	158	0	1
Togo	1.03	0.97	0.98	0.30	0.05	135	140	138	2	3
Trinidad/Tobago	7.56	3.09	6.20	50.15	0.18	5	54	11	43	6
Tunisia	1.76	1.90	1.85	-2.47	-0.05	92	85	85	0	-7
Turkey	2.55	2.70	2.65	-1.88	-0.04	72	66	65	1	-7
Turkmenistan	3.98	3.93	3.88	-1.15	0.03	45	47	45	2	0
Uganda	1.57	1.53	1.60	4.43	-0.02	101	104	99	5	-2
Ukraine	3.19	2.90	2.70	-7.45	0.15	54	60	61	-1	7
U.A.E.	8.44	10.68	10.32	-3.49	-0.22	3	1	2	-1	-1
United Kingdom	4.71	4.89	4.81	-1.78	-0.02	29	33	30	3	1
Tanzania	1.19	1.18	1.23	4.59	-0.04	124	128	121	7	-3
United States	7.19	8.00	7.58	-5.46	-0.05	6	6	4	2	-2
Uruguay	5.08	5.13	5.20	1.33	-0.02	22	27	25	2	3
Uzbekistan	1.82	1.74	1.71	-1.88	0.06	87	96	94	2	7
Venezuela	3.02	2.89	2.72	-6.23	0.10	56	61	59	2	3
Viet Nam	1.39	1.40	1.36	-2.92	0.02	113	111	110	1	-3
Yemen	0.87	0.94	0.96	1.87	-0.10	141	143	140	3	-1
Zambia	0.84	0.91	0.83	-9.27	0.01	145	146	147	-1	2

Zimbabwe 1.17 1.25 1.31 5.03 -0.12 125 121 115 6 -10

629

630

631

TABLE 3: NFA 2010 and NFA 2011 Editions comparison: biocapacity data table.

	(a) Biocapacity 2008 (2011 Edition)	(b) Biocapacity 2007 (2010 Edition)	(c) Biocapacity 2007 (2011 Edition)	(c)- (b) (c)	(a)- (c) (a)	Rank (a)	Rank (b)	Rank (c)	ΔRank between editions - (b) to (c)	ΔRank between years - (c) to (a)
	[gha/capita]	[gha/capita]	[gha/capita]	%	%					
Afghanistan	0.40	0.54	0.47	-15.49	-0.17	140	139	142	-3	2
Albania	0.88	0.87	0.85	-3.08	0.04	107	110	108	2	1
Algeria	0.56	0.59	0.57	-4.15	-0.01	134	138	136	2	2
Angola	2.98	3.00	3.00	0.13	-0.01	39	40	40	0	1
Argentina	7.12	7.50	7.32	-2.35	-0.03	16	15	16	-1	0
Armenia	0.72	0.71	0.74	3.91	-0.03	120	117	114	3	-6
Australia	14.57	14.71	14.54	-1.22	0.00	5	5	5	0	0
Austria	3.34	3.31	3.25	-2.10	0.03	31	33	33	0	2
Azerbaijan	0.72	0.76	0.73	-5.01	-0.01	116	118	116	2	0
Bahrain	0.69	0.94	0.77	-22.04	-0.11	105	114	121	-7	16
Bangladesh	0.42	0.38	0.40	6.91	0.04	150	148	147	1	-3
Belarus	3.40	3.29	3.20	-2.83	0.06	32	35	32	3	0
Belgium	1.33	1.34	1.34	-0.14	-0.01	81	81	84	-3	3
Benin	0.98	0.78	0.98	20.69	0.00	115	103	104	-1	-11
Bolivia	18.39	18.84	18.74	-0.51	-0.02	2	2	2	0	0
Bosnia/Herzegovina	1.64	1.60	1.57	-1.82	0.04	72	74	71	3	-1
Botswana	3.76	3.83	3.79	-0.91	-0.01	28	27	26	1	-2
Brazil	9.63	8.98	9.67	7.13	0.00	12	11	11	0	-1
Bulgaria	2.65	2.13	2.05	-3.84	0.23	59	61	49	12	-10
Burkina Faso	1.37	1.30	1.19	-8.95	0.13	84	90	81	9	-3
Burundi	0.45	0.50	0.48	-4.45	-0.08	141	144	145	-1	4
Cambodia	1.01	0.94	0.97	3.19	0.04	104	104	103	1	-1
Cameroon	1.87	1.85	1.92	3.63	-0.03	66	64	67	-3	1
Canada	14.92	14.92	14.74	-1.21	0.01	4	4	4	0	0
C. African Republic	8.35	8.44	8.51	0.84	-0.02	14	14	14	0	0
Chad	3.17	3.17	3.21	1.25	-0.01	35	34	35	-1	0
Chile	3.74	3.83	3.78	-1.51	-0.01	27	28	27	1	0
China	0.87	0.98	0.86	-14.33	0.02	103	109	109	0	6
Colombia	3.89	3.98	3.95	-0.86	-0.01	24	23	23	0	-1
Congo	12.20	13.27	12.54	-5.77	-0.03	6	6	6	0	0
Costa Rica	1.60	1.90	1.68	-13.01	-0.05	64	71	72	-1	8
Côte d'Ivoire	1.85	1.67	1.79	6.53	0.03	69	67	68	-1	-1
Croatia	2.92	2.50	2.59	3.31	0.11	50	48	43	5	-7
Cuba	0.71	0.74	0.71	-3.80	-0.01	119	120	119	1	0
Cyprus	0.24	0.40	0.29	-39.00	-0.17	147	154	153	1	6
Czech Republic	2.68	2.67	2.60	-2.44	0.03	46	46	47	-1	1
Korea, DPR	0.62	0.58	0.60	3.15	0.04	135	134	130	4	-5
Congo, DR	3.10	2.76	3.19	13.53	-0.03	44	36	36	0	-8
Denmark	4.81	4.85	4.72	-2.75	0.02	21	21	21	0	0
Dominican Rep.	0.54	0.50	0.50	0.58	0.07	142	142	139	3	-3
Ecuador	2.18	2.33	2.22	-4.98	-0.02	54	57	62	-5	8
Egypt	0.65	0.62	0.66	6.71	-0.01	127	125	125	0	-2
El Salvador	0.62	0.67	0.62	-7.95	0.00	123	129	129	0	6
Eritrea	1.47	1.60	1.72	7.09	-0.17	73	70	76	-6	3
Estonia	8.73	8.96	8.95	-0.10	-0.02	13	13	13	0	0
Ethiopia	0.65	0.66	0.65	-2.06	0.00	125	128	127	1	2
Finland	12.19	12.46	12.33	-1.10	-0.01	7	7	7	0	0
France	2.99	3.00	2.92	-2.75	0.02	38	42	39	3	1
Gabon	28.72	29.29	29.24	-0.19	-0.02	1	1	1	0	0
Gambia	1.15	1.10	1.05	-5.13	0.09	98	100	95	5	-3
Georgia	1.17	1.21	1.18	-1.99	-0.01	88	91	94	-3	6
Germany	1.95	1.92	1.88	-2.43	0.04	63	66	66	0	3
Ghana	1.28	1.19	1.26	5.25	0.02	91	86	90	-4	-1
Greece	1.59	1.62	1.50	-7.83	0.05	70	76	73	3	3

Guatemala	1.07	1.12	1.16	3.37	-0.09	96	93	100	-7	4
Guinea	2.93	2.85	2.91	2.25	0.00	40	43	42	1	2
Guinea-Bissau	3.40	3.22	3.49	7.79	-0.03	33	30	31	-1	-2
Haiti	0.31	0.31	0.33	7.94	-0.08	153	150	151	-1	-2
Honduras	1.97	1.84	2.04	9.92	-0.03	67	62	65	-3	-2
Hungary	2.68	2.23	2.15	-3.94	0.20	57	60	46	14	-11
India	0.48	0.51	0.49	-4.61	-0.01	140	143	143	0	3
Indonesia	1.32	1.35	1.32	-2.77	0.00	79	84	87	-3	8
Iran, Islamic Rep.	0.84	0.81	0.92	11.96	-0.10	114	106	111	-5	-3
Iraq	0.24	0.30	0.33	9.37	-0.38	154	151	154	-3	0
Ireland	3.41	3.48	3.49	0.31	-0.02	30	31	30	1	0
Israel	0.29	0.32	0.31	-2.21	-0.07	152	153	152	1	0
Italy	1.15	1.14	1.13	-1.20	0.01	94	95	96	-1	2
Jamaica	0.33	0.38	0.33	-15.61	-0.02	149	152	150	2	1
Japan	0.59	0.60	0.59	-1.62	0.00	131	135	134	1	3
Jordan	0.24	0.24	0.24	-0.08	-0.02	155	155	155	0	0
Kazakhstan	3.48	4.01	3.93	-1.96	-0.13	23	24	29	-5	6
Kenya	0.53	0.59	0.59	-1.17	-0.11	133	136	141	-5	8
Kuwait	0.43	0.40	0.45	12.15	-0.05	148	145	146	-1	-2
Kyrgyzstan	1.33	1.34	1.38	2.43	-0.04	80	80	86	-6	6
Laos	1.65	1.58	1.60	0.94	0.03	74	72	70	2	-4
Latvia	6.63	7.07	6.58	-7.43	0.01	17	17	17	0	0
Lebanon	0.39	0.40	0.38	-5.25	0.03	146	149	149	0	3
Lesotho	0.81	0.81	0.82	0.46	-0.01	113	112	112	0	-1
Liberia	2.95	2.47	3.11	20.41	-0.05	52	37	41	-4	-11
Libya	0.66	0.44	0.68	34.48	-0.02	144	123	124	-1	-20
Lithuania	4.32	4.36	4.24	-2.92	0.02	22	22	22	0	0
Madagascar	2.92	3.07	2.98	-2.94	-0.02	37	41	44	-3	7
Malawi	0.67	0.70	0.72	2.15	-0.08	121	119	123	-4	2
Malaysia	2.50	2.61	2.52	-3.33	-0.01	48	49	51	-2	3
Mali	2.29	2.49	2.19	-13.68	0.04	51	59	57	2	6
Mauritania	5.21	5.50	5.34	-2.96	-0.02	19	20	20	0	1
Mauritius	0.56	0.56	0.55	-1.88	0.02	138	139	137	2	-1
Mexico	1.42	1.47	1.42	-3.41	0.00	76	78	78	0	2
Mongolia	15.33	15.14	15.61	3.03	-0.02	3	3	3	0	0
Morocco	0.70	0.61	0.61	0.49	0.12	130	132	120	12	-10
Mozambique	2.21	1.89	2.25	15.81	-0.02	65	56	60	-4	-5
Myanmar	2.22	2.04	2.19	6.83	0.01	61	58	59	-1	-2
Namibia	7.18	7.56	7.31	-3.42	-0.02	15	16	15	1	0
Nepal	0.53	0.55	0.53	-3.21	0.01	139	140	140	0	1
Netherlands	1.03	1.03	1.01	-1.74	0.02	99	102	101	1	2
New Zealand	10.19	10.77	10.35	-4.04	-0.02	9	9	9	0	0
Nicaragua	2.33	2.82	2.39	-18.08	-0.03	41	52	53	-1	12
Nigeria	1.12	1.12	1.08	-3.68	0.04	97	99	97	2	0
Norway	5.40	5.48	5.41	-1.33	0.00	20	19	19	0	-1
Palestinian Terr.	0.13	0.16	0.13	-21.13	0.01	156	157	157	0	1
Oman	2.20	2.14	2.26	5.21	-0.03	58	55	61	-6	3
Pakistan	0.40	0.43	0.44	1.68	-0.08	145	146	148	-2	3
Panama	2.67	3.15	2.70	-16.59	-0.01	36	44	48	-4	12
Papua New Guinea	3.67	3.75	3.73	-0.63	-0.02	29	29	28	1	-1
Paraguay	10.92	11.24	11.07	-1.54	-0.01	8	8	8	0	0
Peru	3.82	3.86	3.85	-0.33	-0.01	26	26	25	1	-1
Philippines	0.62	0.62	0.61	-2.00	0.02	126	133	131	2	5
Poland	2.00	2.09	2.03	-2.94	-0.01	60	63	64	-1	4
Portugal	1.29	1.25	1.27	1.20	0.02	85	85	89	-4	4
Puerto Rico	0.17	0.14	0.17	16.92	-0.05	157	156	156	0	-1
Qatar	2.05	2.51	2.42	-3.88	-0.18	49	51	63	-12	14
Korea, Rep.	0.72	0.33	0.70	52.47	0.02	151	122	115	7	-36
Moldova	1.33	0.66	0.65	-1.79	0.51	124	126	85	41	-39
Romania	2.33	1.95	1.88	-3.71	0.19	62	65	54	11	-8
Russia	6.62	5.75	6.52	11.78	0.02	18	18	18	0	0
Rwanda	0.52	0.56	0.52	-9.03	0.01	136	141	142	-1	6
Saudi Arabia	0.65	0.84	0.68	-23.97	-0.03	112	124	126	-2	14
Senegal	1.40	1.20	1.23	2.40	0.12	89	87	80	7	-9
Serbia	1.41	1.16	1.20	3.22	0.15	92	89	79	10	-13
Sierra Leone	1.71	1.20	1.73	30.81	-0.01	90	69	69	0	-21
Singapore	0.02	0.02	0.02	-2.19	-0.04	158	158	158	0	0
Slovakia	2.86	2.68	2.61	-2.41	0.09	45	45	45	0	0

Slovenia	2.59	2.61	2.60	-0.23	0.00	47	47	50	-3	3
Somalia	1.36	1.40	1.41	1.21	-0.04	78	79	82	-3	4
South Africa	1.21	1.14	1.14	0.17	0.05	95	94	91	3	-4
Spain	1.46	1.61	1.58	-1.94	-0.09	71	73	77	-4	6
Sri Lanka	0.46	0.45	0.43	-4.04	0.07	143	147	144	3	1
Sudan	2.34	2.42	2.45	1.03	-0.05	53	50	52	-2	-1
Swaziland	0.97	1.00	0.96	-4.25	0.00	101	105	105	0	4
Sweden	9.51	9.75	9.67	-0.79	-0.02	11	12	12	0	1
Switzerland	1.20	1.24	1.22	-1.88	-0.02	86	88	92	-4	6
Syria	0.57	0.70	0.71	0.98	-0.24	122	121	135	-14	13
Tajikistan	0.56	0.56	0.58	4.10	-0.05	137	137	138	-1	1
Thailand	1.17	1.15	1.17	1.08	0.01	93	92	93	-1	0
Macedonia TFYR	1.55	1.43	1.47	2.23	0.06	77	77	75	2	-2
Timor-Leste	0.86	1.21	0.88	-38.22	-0.02	87	108	110	-2	23
Togo	0.67	0.60	0.65	8.13	0.03	132	127	122	5	-10
Trinidad/Tobago	1.56	1.57	1.57	0.42	-0.01	75	75	74	1	-1
Tunisia	0.96	0.98	1.01	3.24	-0.06	102	101	106	-5	4
Turkey	1.31	1.32	1.33	0.76	-0.02	83	82	88	-6	5
Turkmenistan	3.19	3.21	3.30	2.63	-0.03	34	32	34	-2	0
Uganda	0.81	0.85	0.81	-4.14	0.00	111	113	113	0	2
Ukraine	2.23	1.82	1.77	-2.96	0.21	68	68	58	10	-10
U.A.E.	0.64	0.85	0.83	-1.85	-0.30	110	111	128	-17	18
United Kingdom	1.34	1.34	1.32	-1.58	0.02	82	83	83	0	1
Tanzania	1.02	1.02	1.08	5.98	-0.06	100	98	102	-4	2
United States	3.86	3.87	3.87	-0.03	0.00	25	25	24	1	-1
Uruguay	10.03	9.91	9.91	-0.01	0.01	10	10	10	0	0
Uzbekistan	0.91	0.92	0.91	-1.11	0.00	106	107	107	0	1
Venezuela	3.00	2.81	3.06	7.95	-0.02	42	38	37	1	-5
Viet Nam	1.09	0.86	1.09	21.12	0.00	108	96	98	-2	-10
Yemen	0.60	0.62	0.62	0.20	-0.04	128	130	132	-2	4
Zambia	2.31	2.26	2.38	4.98	-0.03	55	53	55	-2	0
Zimbabwe	0.72	0.75	0.76	1.03	-0.06	117	115	117	-2	0

632

633

634 Methodological differences between editions can be demonstrated by looking at the change in absolute  
635 Ecological Footprint and biocapacity, and by looking at changes in country rankings for these two  
636 indicators. For the year 2007 - the most recent year covered by both NFA 2011 and NFA 2010 Editions  
637 - there were seven countries whose rank in Ecological Footprint per capita changed more than 15 places  
638 (standard deviation - s.d. = 12.1); for biocapacity per capita, there were only two countries whose rank  
639 changed by more than 15 places (s.d. = 5.2). Nine countries showed absolute changes in the Ecological  
640 Footprint greater than 1.0 gha per capita (s.d. = 0.6 gha per capita); no countries showed absolute  
641 changes in biocapacity greater than 1.0 gha per capita (s.d. = 0.2 gha per capita) (Figure 6).

642

643

644

645

646

647


648

649


650

651

652 FIGURE 6: Histogram of changes in country ranks (top) and per capita values (bottom) moving from  
 653 NFA 2010 edition to NFA 2011 edition for the data year 2007. Lines indicate normal distribution fit to  
 654 each histogram.



655



656

657


658 Regardless of the changes at the national level, trends for both editions show an overall decrease in  
 659 world biocapacity and an overall increase in Ecological Footprint during the 47 year time series. Figure  
 660 7 shows the trend for humanity's average per capita Ecological Footprint and biocapacity for both the  
 661 2010 and 2011 Editions of the National Footprint Accounts. The largest difference between the two  
 662 editions is the increasing difference in biocapacity going back in time, obtained as a result of the shift to  
 663 a constant global hectare approach (see section 5.2). Due to the increase in agricultural productivity  
 664 over the last 50 years, one hectare of cropland in 1961 provided fewer resources for human

665 consumption than one hectare of cropland in 2008, and thus corresponds to fewer constant global  
666 hectares of biocapacity.

667

668 FIGURE 7: Humanity's average per capita Ecological Footprint (EF) and biocapacity (BC) over time.  
669 Trends from the 2008, 2009, 2010 and 2011 Editions of the National Footprint Accounts are provided  
670 for comparison purposes.

671



672

673

674 A similar reduction in the Ecological Footprint takes place when moving to a constant global hectare calculation.  
675 However, this change has been largely offset by the change in the ocean uptake calculation (see section 5.3),  
676 where the NFA 2011 Edition uses a much lower value of ocean sequestration than prior versions, and thus there  
677 is an increased carbon Ecological Footprint. Taken together, these two methodological changes result in a large  
678 shift in the relative composition of the 1961 Ecological Footprint between NFA 2010 and NFA 2011 (48%  
679 cropland/12% carbon and 24% cropland /36% carbon respectively).

680

681 Nevertheless, global trends in the Ecological Footprint and biocapacity show a consistent message  
682 across the last four methodological updates of the National Footprint Accounts: population growth that  
683 outstrips increases in bioproductivity; and, following a relatively rapid increase in the 1960s, little  
684 change in the average Ecological Footprint per person over the last 40 years.

685

## 686 7. National Footprint Accounts' limitations

687 NFAs aim at measuring whether or not humans are able to live within the Biosphere's ecological  
688 budget. To answer this research question, a systemic approach is used to assess, in a combined way, the  
689 impact of pressures that are usually evaluated independently. Therefore, NFAs have been developed as

690 a resource accounting framework, where the various pressures are first analyzed independently and  
691 results are then aggregated into a single number (see section 2 and Figure 1). Aggregation, however, has  
692 the drawback of implying a greater degree of additivity and substitutability between the included land  
693 use types than is probably realistic (DG Environment, 2008; Giljum et al., 2009; Kitzes et al., 2009;  
694 Wiedmann and Barrett, 2010).

695

696 The quality, reliability and validity of the NFAs are dependent upon the level of accuracy and  
697 availability of a wide range of datasets, many of which have incomplete coverage, and most of which  
698 do not specify confidence limits. Considerable care is taken to minimize any data inaccuracies or  
699 calculation errors that might distort the NFAs, including inviting national governments to  
700 collaboratively review the assessment of their country for accuracy (e.g., Abdullatif and Alam, 2011;  
701 Hild et al., 2010; von Stokar et al., 2006). In addition, the Ecological Footprint methodology is  
702 continually being refined and efforts are made to improve the transparency of the NFAs and the related  
703 written documentation (Gracey et al., 2012; Kitzes et al., 2009), allowing for more effective internal  
704 and external review.

705

706 Finally, NFAs are specifically constructed to yield conservative estimates of global overshoot. On the  
707 supply side, biocapacity is overestimated as both the land degradation and the long-term sustainability  
708 of resource extraction is not taken into account. On the supply side, Ecological Footprint is  
709 underestimated as it does not track freshwater consumption, soil erosion, GHGs emissions other than  
710 CO<sub>2</sub> as well as impacts for which no regenerative capacity exists (e.g. pollution in terms of waste  
711 generation, toxicity, eutrophication, etc). A detailed list of strengths and weaknesses of the Ecological  
712 Footprint methodology and limitations of the NFAs, can be found in Galli et al (2011) and Ewing et al  
713 (2010b), respectively.

714

## 715 **Conclusions**

716 In an increasingly resource constrained world, accurate and effective resource accounting systems are  
717 needed if nations, cities and companies want to stay competitive. National Footprint Accounts (NFA) is  
718 one such accounting system, designed to track human demand on the regenerative and absorptive  
719 capacity of the biosphere.

720

721 NFAs are maintained and updated annually by Global Footprint Network. Every new edition relies on  
722 the use of more comprehensive data sets and independent data sources, more consistent and reliable  
723 data, a revised and updated methodology and a more robust calculation process. Each time a new



724 edition is released, Ecological Footprint and biocapacity values are back calculated from the most  
725 recent year in order to ensure consistency across the historical time trends. Edition after edition, these  
726 improvements lead to more reliable (and yet consistent) Ecological Footprint and biocapacity values  
727 and trends for nations and the world.

728

729 Stakeholders interested in monitoring nations' Ecological Footprint and biocapacity values and/or  
730 setting Footprint reduction targets are advised not to compare results obtained via different editions of  
731 the NFAs, and encouraged to always look at the time trends from the most recent edition of the NFAs.

732 **Acknowledgments**

733 We would like to thank Robert Williams for his help in the National Footprint Accounts Program as  
734 well as Nicolò Passeri for comments and feedback on a preliminary version of the paper. Thanks are  
735 also due to the two anonymous reviewers for their precious and constructive comments.

736

737 We would also like to acknowledge Avina Stiftung, Arthur und Estella Hirzel-Callegari Stiftung,  
738 Environment Agency - Abu Dhabi, Flora Family Foundation, Foundation for Global Community, Karl  
739 Mayer Stiftung, MAVA - Fondation pour la Protection de la Nature, Mental Insight Foundation, Skoll  
740 Foundation, Stiftung ProCare, Winslow Foundation, WWF International, Zayed International Prize for  
741 the Environments and many individual donors who have made the update of the National Footprint  
742 accounts possible. Thanks also go to the National Footprint Accounts Review Committee who has been  
743 providing continuous feedback on the development of the methodology.

744

745 **References**

746

747 Abdullatif, L., Alam, T., 2011. *The UAE Ecological Footprint Initiative*.

748 [http://awsassets.panda.org/downloads/en\\_final\\_report\\_ecological\\_footprint.pdf](http://awsassets.panda.org/downloads/en_final_report_ecological_footprint.pdf)

749 Akaike, H., 1978. A Bayesian Analysis of the Minimum AIC Procedure. *Ann.Inst. Statist. Math.* 30,

750 Part A, 9-14.

751 Barnosky AD, Hadly EA, Bascompte J, Berlow EL, Brown JH, Fortelius M, Getz WM, Harte J,  
752 Hastings A, Marquet PA, Martinez ND, Mooers A, Roopnarine P, Vermeij G, Williams JW,  
753 Gillespie R, Kitzes J, Marshall C, Matzke N, Mindell DP, Revilla E, Smith AB. 2012.

754 Approaching a state shift in Earth's biosphere. *Nature* 486, 52-58.

755 Bastianoni, S., Niccolucci, V., Pulselli, R.M., Marchettini, N., 2012. Indicator and *indicandum*:

756 "Sustainable way" vs "prevailing conditions" in the Ecological Footprint. *Ecological Indicators*,

757 16, 47–50.

758 Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S.,

759 O'Neill, R.V., Paruelo, J., Raskin, R.G., Sutton, P., van den Belt, M. 1997. The value of the

760 world's ecosystem services and natural capital. *Nature*, 387, 253-260.

761 Costanza, R., Daly, H.E., 1992. Natural capital and sustainable development. *Conservation Biology* 6,

762 37–46.

763 Daly, H.E., Farley, J., 2004. *Ecological Economics: Principles and Application*. Island Press,

764 Washington, USA.

765 Daly, H.E., 1990. Towards Some Operational Principles of Sustainable Development. *Ecological*

766 *Economics* 2, 1–6.

767 DeFries, R.S., Foley, J.A., Asner, G.P., 2004. Land-use choices: balancing human needs and ecosystem

768 function. *Front Ecol Environ* 2(5), 249–257.

769 DG Environment, 2008. *Potential of the Ecological Footprint for monitoring environmental impact*

770 *from natural resource use*. <http://ec.europa.eu/environment/natres/studies.htm>.

771 Ewing, B., Reed, A., Galli, A., Kitzes, J., Wackernagel, M., 2010a. *Calculation Methodology for the*

772 *national Footprint Accounts, 2010 Edition*. Oakland: Global Footprint Network

773 [http://www.footprintnetwork.org/images/uploads/National\\_Footprint\\_Accounts\\_Method\\_Paper](http://www.footprintnetwork.org/images/uploads/National_Footprint_Accounts_Method_Paper)

774 [2010.pdf](http://www.footprintnetwork.org/images/uploads/National_Footprint_Accounts_Method_Paper_2010.pdf)

775 Ewing, B., Moore, D., Goldfinger, S., Oursler, A., Reed, A., Wackernagel, M., 2010b. *The Ecological*

776 *Footprint Atlas 2010*. Oakland: Global Footprint Network. [www.footprintnetwork.org/atlas](http://www.footprintnetwork.org/atlas).

777 Food and Agriculture Organization of the United Nations (FAO) Statistical Databases.

778 <http://faostat.fao.org/site/291/default.aspx> (accessed February 2011).

779 FAO, 2000. *Technical Conversion Factors for Agricultural Commodities*.

780 <http://www.fao.org/es/ess/tcf.asp>. (accessed February 2011).

781 FAO, 2006. *World maps of climatological net primary production of biomass, NPP*.

782 [http://www.fao.org/nr/climpag/globgrids/NPP\\_en.asp](http://www.fao.org/nr/climpag/globgrids/NPP_en.asp) (accessed March 2012)

- 783 FAO and IIASA (International Institute for Applied Systems Analysis), 2000. Global Agro-Ecological  
784 Zones.. <http://www.fao.org/ag/agl/agll/gaez/index.htm> (accessed February 2011).
- 785 FAO, 1998. *Global Fiber Supply Model*. <ftp://ftp.fao.org/docrep/fao/006/X0105E/X0105E.pdf>  
786 (accessed February 2011).
- 787 Ferguson, A.R.B., 1999. The logical foundations of ecological footprints. *Environment, Development*  
788 *and Sustainability* 1, 149–156.
- 789 Galli, A., 2007. *Assessing the role of the Ecological Footprint as Sustainability Indicator*. Ph.D. Thesis.  
790 Department of Chemical and Biosystems Sciences. University of Siena, Italy.
- 791 Galli, A., Kitzes, J., Wermer, P., Wackernagel, M., Niccolucci, V., Tiezzi, E., 2007. An Exploration of  
792 the Mathematics behind the Ecological Footprint. *International Journal of Ecodynamics*. 2(4),  
793 250-257.
- 794 Galli, A., Wiedmann, T., Ercin, E., Knoblauch, D., Ewing, B., Giljum, S., 2012. Integrating Ecological,  
795 Carbon, and Water Footprint into a “Footprint Family” of indicators: definition and role in  
796 tracking Human Pressure on the Planet. *Ecological Indicators*, 16, 100-112.
- 797 Galli, A., Wiedmann, T., Ercin, E., Knoblauch, D., Ewing, B., Giljum, S., 2011. *Integrating Ecological,*  
798 *Carbon and Water Footprint: Defining the “Footprint Family” and its Application in Tracking*  
799 *Human Pressure on the Planet*. OPEN:EU project deliverable.  
800 [http://www.oneplaneteconomynetwork.org/resources/programme-  
801 documents/WP8\\_Integrating\\_Ecological\\_Carbon\\_Water\\_Footprint.pdf](http://www.oneplaneteconomynetwork.org/resources/programme-<br/>801 documents/WP8_Integrating_Ecological_Carbon_Water_Footprint.pdf)
- 802 Giljum, S., Hinterberger, F., Lutter, S., Polzin, C., 2009. *How to Measure Europe’s Resource Use. An*  
803 *Analysis for Friends of the Earth Europe*. Sustainable Europe Research Institute, Vienna.
- 804 Global Footprint Network, 2011. *National Footprint Accounts, 2011 Edition*. Available at  
805 [www.footprintnetwork.org](http://www.footprintnetwork.org).
- 806 Gracey, K., Lazarus, E., Borucke, M., Moore, D., Cranston, G., Iha, K., Larson, J., Morales, J.C.,  
807 Wackernagel, M., Galli, A., 2012. *Guidebook to the National Footprint Accounts: 2011 Edition*.  
808 Oakland: Global Footprint Network. [www.footprintnetwork.org/methodology](http://www.footprintnetwork.org/methodology).
- 809 Gulland, J.A., 1971. *The Fish Resources of the Ocean*. West Byfleet, Surrey, United Kingdom: Fishing  
810 News.
- 811 Haberl, H., Erb, K., Krausmann, F., 2001. How to calculate and interpret ecological footprints for long  
812 periods of time: the case of Austria 1926-1995. *Ecological Economics*, 38, 25-45.
- 813 Haberl, H., Erb, K.H., Krausmann, F., Gaube, V., Bondeau, A., Plutzer, C., Gingrich, S., Lucht, W.,  
814 Fischer-Kowalski, M., 2007. Quantifying and mapping the human appropriation of net primary  
815 production in earth’s terrestrial ecosystems. *Proc. Natl. Acad. Sci.* 104, 12942-12947.
- 816 Hammond, G.P., Jones, C.I., 2008. Embodied energy and carbon in construction materials. *Proc. Instn*  
817 *Civil. Engrs: Energy*, 161(2), 87-98.
- 818 Heinberg, R., 2007. *Peak Everything: Waking Up to the Century of Declines*. New Society Publishers,  
819 Gabriola Island, British Columbia, Canada.

- 820 Hild, P., Schmitt, B., Decoville, A., Mey, M., Welfring, J., 2010. *The Ecological Footprint of*  
821 *Luxembourg - Technical Report*. [http://www.myfootprint.lu/files/RAP-20100614-](http://www.myfootprint.lu/files/RAP-20100614-EF_Lux_Phase_2-v.4.0.pdf)  
822 [EF\\_Lux\\_Phase\\_2-v.4.0.pdf](http://www.myfootprint.lu/files/RAP-20100614-EF_Lux_Phase_2-v.4.0.pdf)
- 823 IEA Statistics and Balances. <http://data.iea.org/ieastore/statslisting.asp> (accessed February 2011).
- 824 Imhoff, M.L., Lawrence, W.T., Elvidge, C., Paul, T., Levine, E., Privalsky, M., Brown, V., 1997. Using  
825 nighttime DMSP/OLS images of city lights to estimate the impact of urban land use on soil  
826 resources in the United States. *Remote Sens. Environ.* 59, 105–117.
- 827 Interfacultaire Vakgroep Energie en Milieukunde Energy Analysis Program, Research Report no. 98,  
828 Groningen, 1999.
- 829 IPCC (Intergovernmental Panel on Climate Change), 2001. *Climate Change 2001: The Scientific Basis*.  
830 Cambridge, UK: Cambridge University Press, 2001.
- 831 IPCC (Intergovernmental Panel on Climate Change), 2006. *2006 IPCC Guidelines for National*  
832 *Greenhouse Gas Inventories Volume 4: Agriculture Forestry and Other Land Use*.  
833 <http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol4.html> (accessed February 2010).  
834
- 835 IPCC (Intergovernmental Panel on Climate Change), 2007. *Climate Change 2007: The Physical*  
836 *Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the*  
837 *Intergovernmental Panel on Climate Change*. [Solomon, S., D. Qin, M. Manning, Z. Chen, M.  
838 Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press,  
839 Cambridge, United Kingdom and New York, NY, USA.
- 840 Khatiwala, S., Primeau, F., Hall, T., 2009. Reconstruction of the history of anthropogenic CO<sub>2</sub>  
841 concentrations in the ocean. *Nature*, 462, 346-350.
- 842 Kitzes, J., Galli, A., Bagliani, M., Barrett, J., Dige, G., Ede, S., Erb, K-H., Giljum, S., Haberl, H., Hails,  
843 C., Jungwirth, S., Lenzen, M., Lewis, K., Loh, J., Marchettini, N., Messinger, H., Milne, K.,  
844 Moles, R., Monfreda, C., Moran, D., Nakano, K., Pyhälä, A., Rees, W., Simmons, C.,  
845 Wackernagel, M., Wada, Y., Walsh, C., Wiedmann, T., 2009. A research agenda for improving  
846 national ecological footprint accounts. *Ecological Economics*, 68(7), 1991-2007.
- 847 Kitzes, J., Peller, A., Goldfinger, S., Wackernagel, M., 2007a. Current Methods for Calculating  
848 National Ecological Footprint Accounts. *Science for Environment & Sustainable Society*, 4(1)  
849 1-9.
- 850 Kitzes, J., Galli, A., Wackernagel, M., Goldfinger, S., Bastianoni, S., 2007b. A ‘constant global hectare’  
851 method for representing ecological footprint time trends. Available on-line [URL]:  
852 [http://www.brass.cf.ac.uk/uploads/fullpapers/Kitzes\\_M66.pdf](http://www.brass.cf.ac.uk/uploads/fullpapers/Kitzes_M66.pdf).
- 853 Marland, G., Boden, T.A., Andres, R.J., 2007. *Global, Regional, and National Fossil Fuel CO<sub>2</sub>*  
854 *Emissions*. In Trends: A Compendium of Data on Global Change. Oak Ridge, TN: Carbon  
855 Dioxide Information Analysis Center, Oak Ridge National Laboratory and U.S. Department of  
856 Energy.
- 857 Max-Neef, M., 1995. Economic growth and quality of life: a threshold hypothesis. *Ecological*  
858 *Economics* 15, 115–118.
- 859 Monfreda, C., Wackernagel, M., Deumling, D., 2004. Establishing national natural capital accounts  
860 based on detailed ecological footprint and biocapacity assessments. *Land Use Policy* 21, 231–  
861 246.

- 862 Moore, D., Galli, A., Cranston, G.R., Reed, A. Projecting future human demand on the Earth's  
863 regenerative capacity. *Ecological Indicators* 16, 3-10.
- 864 Pauly, D., Christensen, V., 1995. Primary production required to sustain global fisheries. *Nature*, 374,  
865 255-257. Rees, W.E., 1992. Ecological footprints and appropriated carrying capacity: What  
866 urban economics leaves out. *Environment and Urbanization*, 4, 121-130.
- 867 Rockström, J., W. Steffen, K. Noone, Å. Persson, F. S. Chapin, III, E. Lambin, T. M. Lenton, M.  
868 Scheffer, C. Folke, H. Schellnhuber, B. Nykvist, C. A. De Wit, T. Hughes, S. van der Leeuw, H.  
869 Rodhe, S. Sörlin, P. K. Snyder, R. Costanza, U. Svedin, M. Falkenmark, L. Karlberg, R. W.  
870 Corell, V. J. Fabry, J. Hansen, B. Walker, D. Liverman, K. Richardson, P. Crutzen, and J. Foley.  
871 2009a. Planetary boundaries: exploring the safe operating space for humanity. *Ecology and*  
872 *Society* 14(2): 32. [online] URL: <http://www.ecologyandsociety.org/vol14/iss2/art32/>
- 873 Rockström R, Steffen W, Noone K, Persson A, Chapin FS, Lambin EF, Lenton TM, Scheffer M, Folke  
874 C, Schellnhuber HJ, Nykvist B, de Wit CA, Hughes T, van der Leeuw S, Rodhe H, Sörlin S,  
875 Snyder PK, Costanza R, Svedin U, Falkenmark M, Karlberg L, Corell RW, Fabry VJ, Hansen J,  
876 Walker B, Liverman D, Richardson K, Crutzen P, Foley JA., 2009b. A safe operating space for  
877 humanity. *Science*, 46, 472-475.
- 878 Scheffer, M., Carpenter, S., Foley, J.A., Folke, C., Walker, B., 2001. Catastrophic shifts in ecosystems.  
879 *Nature* 413, 591-596.
- 880 Schlesinger, W.H. 2009. Planetary boundaries: Thresholds risk prolonged degradation. *Nature Reports*  
881 *Climate Change* 3, 112-113.
- 882 Stiglitz, J.E., Sen, A., Fitoussi, J-P., 2009. *Report by the Commission on the Measurement of Economic*  
883 *Performance and Social Progress*. [http://www.stiglitz-sen-](http://www.stiglitz-sen-fitoussi.fr/documents/rapport_anglais.pdf)  
884 [fitoussi.fr/documents/rapport\\_anglais.pdf](http://www.stiglitz-sen-fitoussi.fr/documents/rapport_anglais.pdf)
- 885 Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C., Erasmus,  
886 B.F.N., Ferreira de Siqueira, M., Grainger, A., Hannah, L., Hughes, L., Huntley, B., van  
887 Jaarsveld, A.S., Midgley, G.F., Miles, L., Ortega-Huerta, M.A., Peterson, A.T., Phillips, O.L.,  
888 Williams, S.E., 2004. Extinction risk from climate change. *Nature* 427, 145-148.
- 889 Thormark, C., 2002. A low energy building in a life cycle-its embodied energy, energy need for  
890 operation and recycling potential. *Building and Environment*, 37, 429-435.
- 891 UN Commodity Trade Statistics Database. 2011. <http://comtrade.un.org> (accessed February 2011).
- 892 UNECE and FAO, 2000. *Temperate and Boreal Forest Resource Assessment*. Geneva: UNECE, FAO.
- 893 UNEP (United Nations Environment Programme), 2007. *GEO4 Global Environment Outlook:*  
894 *environment for development*. Progress Press Ltd, Malta.
- 895 Venetoulis, J., Talberth, J., 2008. Refining the ecological footprint. *Environment, Development and*  
896 *Sustainability*, 10(4), 441-469.
- 897 von Stokar, T., Steinemann, M., Rüegge, B., 2006. *Ecological Footprint of Switzerland - Technical*  
898 *Report*. Published by Federal Office for Spatial Development (ARE), Agency for Development and  
899 Cooperation (SDC), Federal Office for the Environment (FOEN), and Federal Statistical Office  
900 (FSO).  
901 [http://www.bfs.admin.ch/bfs/portal/en/index/themen/21/03/01\\_parsys.0001\\_downloadList.00011\\_DownloadFile.tmp/ecologicalfootprinttechnicalreport.pdf](http://www.bfs.admin.ch/bfs/portal/en/index/themen/21/03/01_parsys.0001_downloadList.00011_DownloadFile.tmp/ecologicalfootprinttechnicalreport.pdf)  
902  
903
- 904 Wackernagel, M., 1991. "Land Use: Measuring a Community's Appropriated Carrying Capacity as an  
905 Indicator for Sustainability" and "Using Appropriated Carrying Capacity as an Indicator,  
906 Measuring the Sustainability of a Community". Report I and II to the UBC Task Force on Healthy  
907 and Sustainable Communities. The University of British Columbia, Vancouver.

- 908  
 909 Wackernagel, M., 1994. *Ecological Footprint and Appropriated Carrying Capacity: A Tool for*  
 910 *Planning Toward Sustainability*. Ph.D. Thesis. School of Community and Regional Planning.  
 911 The University of British Columbia.
- 912 Wackernagel, M., Onisto, L., Linares, A.C., Falfán, I.S.L., García, J.M., Guerrero, A.I.S., Guerrero,  
 913 M.G.S., 1997. *Ecological Footprints of Nations: How Much Nature Do They Use? How Much*  
 914 *Nature Do They Have?* Commissioned by the Earth Council for the Rio+5 Forum. Distributed  
 915 by the International Council for Local Environmental Initiatives, Toronto.
- 916 Wackernagel, M., Onisto, L., Bello, P., Linares, A.C., Falfán, I.S.L., García, J.M., Guerrero, A.I.S.  
 917 Guerrero, M.G.S., 1999a. National natural capital accounting with the ecological footprint  
 918 concept, *Ecological Economics*, 29, 375-390.
- 919 Wackernagel, M., Lewan, L. Hansson, C.B., 1999b. Evaluating the use of natural capital with the  
 920 ecological footprint. *Ambio*, 28, 604–612.
- 921 Wackernagel, M., Schulz, B., Deumling, D., Linares, A.C., Jenkins, M., Kapos, V., Monfreda, C., Loh,  
 922 J., Myers, N., Norgaard, R., Randers, J., 2002. Tracking the ecological overshoot of the human  
 923 economy, *Proc. Natl. Acad. Sci*, 99(14), 9266-9271.
- 924 Wackernagel, M., Monfreda, C., Schulz, N.B., Erb, K.H., Haberl, H., Krausmann, F., 2004. Calculating  
 925 national and global ecological footprint time series: resolving conceptual challenges. *Land Use*  
 926 *Policy*, 21(3), 271-278.
- 927 Wackernagel, M., Rees, W.E., 1996. *Our Ecological Footprint: Reducing Human Impact on the Earth*.  
 928 New Society Publishers, Gabriola Island, BC.
- 929 Wackernagel, M., Monfreda, C., Moran, D., Wermer, P., Goldfinger, S., Deumling, D., Murray, M.,  
 930 2005. *National Footprint and Biocapacity Accounts 2005: The underlying calculation method*.  
 931 Oakland: Global Footprint Network. [www.footprintnetwork.org](http://www.footprintnetwork.org).
- 932 Wiedmann, T., Barrett, J., 2010. A review of the ecological footprint indicator—perceptions and  
 933 methods. *Sustainability*, 2 (6), 1645-1693